

REPORT DOCUMENTATION PAGE		Form Approved OMB NO. 0704-0188	
Public Reporting Burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comment regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA, 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington DC 20503			
1. AGENCY USE ONLY (Leave Blank)		2. REPORT DATE:	
		3. REPORT TYPE AND DATES COVERED Final Report 6-Dec-2002 - 28-Aug-2005	
4. TITLE AND SUBTITLE Enhancement of the Anthrax AVA Vaccine with CpG ODN's		5. FUNDING NUMBERS DAAD1903C0002	
6. AUTHORS Arthur M. Krieg, Iain Sim		8. PERFORMING ORGANIZATION REPORT NUMBER	
7. PERFORMING ORGANIZATION NAMES AND ADDRESSES Coley Pharmaceutical Group, Inc. 93 Worcester St. Suite 101 Wellesley, MA 02481 -			
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709-2211		10. SPONSORING / MONITORING AGENCY REPORT NUMBER 44498-LS-DRP.1	
11. SUPPLEMENTARY NOTES The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision, unless so designated by other documentation.			
12. DISTRIBUTION AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited		12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) The abstract is below since many authors do not follow the 200 word limit			
14. SUBJECT TERMS vaccine, anthrax, adjuvant, immune activator		15. NUMBER OF PAGES Unknown due to possible attachments	
		16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION ON THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL

Report Title

Final Report on ACCELERATE ANTHRAX: CpG 7909 Vaccine Adjuvant Program

ABSTRACT

A clinical study “Phase 1/2, Proof-of-Concept, Double-Blind, Randomized, Controlled Trial Assessing the Immunogenicity and Safety of Anthrax Vaccine Adsorbed (BioThrax?) Combined with CPG 7909 in Normal Volunteers” was completed and presented at the 2005 Interscience Conference on Antimicrobial Agents and Chemotherapy in a poster entitled “Marked Enhancement Of Antibody Response To Anthrax Vaccine Adsorbed With CPG 7909 In Healthy Volunteers“. The conclusions from this study included:

1. AVA plus CPG 7909 was reasonably well tolerated
2. There was a trend to greater frequency and severity of adverse events in the AVA plus CPG 7909 group compared to the AVA alone and CPG 7909 alone groups but this was not statistically significant
3. AVA plus CPG 7909 elicited a heightened (6 to 8-fold increase) and accelerated (21 to 24 days) antibody response ($p < 0.001$) compared to AVA alone
4. AVA plus CPG 7909 elicited a positive anti-PA antibody response in $>50\%$ of subjects within 14 days of a single immunization

A robust, large scale manufacturing process of CPG 7909 has been developed and is judged suitable for the supply of drug for further clinical studies and product commercialization.

List of papers submitted or published that acknowledge ARO support during this reporting period. List the papers, including journal references, in the following categories:

(a) Papers published in peer-reviewed journals (N/A for none)

Number of Papers published in peer-reviewed journals: 0.00

(b) Papers published in non-peer-reviewed journals or in conference proceedings (N/A for none)

Number of Papers published in non peer-reviewed journals: 0.00

(c) Papers presented at meetings, but not published in conference proceedings (N/A for none)

MARKED ENHANCEMENT OF ANTIBODY RESPONSE TO ANTHRAX VACCINE ADSORBED WITH CPG 7909 IN HEALTHY VOLUNTEERS

D. RYNKIEWICZ¹, M. RATHKOPF², J. RANSOM³, I. SIM⁴, L. GIRI⁵, J. QUINN², T. WAYTES⁵, M. AL-ADHAMI⁴, W. JOHNSON⁶, C. NIELSEN⁷ ¹V.A. & U. of Texas Health Science Center, San Antonio, TX; ²Wilford Hall Medical Ctr, San Antonio, TX; ³Fast Track Drugs & Biologics, Potomac, MD; ⁴Coley Pharmaceutical Group, Wellesley, MA; ⁵Emergent BioSolutions, Gaithersburg, MD; ⁶USAMRIID, Ft. Detrick, MD; ⁷Consultant to DARPA, Arlington, VA

Number of Papers not Published: 1.00

(d) Manuscripts

Number of Manuscripts: 0.00

Number of Inventions:

Graduate Students

NAME

PERCENT SUPPORTED

FTE Equivalent:

Total Number:

Names of Post Doctorates

NAME

PERCENT SUPPORTED

FTE Equivalent:

Total Number:

Names of Faculty Supported

NAME

PERCENT SUPPORTED

FTE Equivalent:

Total Number:

Names of Under Graduate students supported

NAME

PERCENT SUPPORTED

FTE Equivalent:

Total Number:

Names of Personnel receiving masters degrees

NAME

Total Number:

Names of personnel receiving PHDs

NAME

Total Number:

Names of other research staff

NAME

PERCENT SUPPORTED

FTE Equivalent:

Total Number:

Sub Contractors (DD882)

Inventions (DD882)

MARKED ENHANCEMENT OF ANTIBODY RESPONSE TO ANTHRAX VACCINE ADSORBED WITH CPG 7909 IN HEALTHY VOLUNTEERS


D. RYNKIEWICZ¹, M. RATHKOPF², J. RANSOM³, I. SIM⁴, L. GIRI⁵, J. QUINN², T.
WAYTES⁵, M. AL-ADHAMI⁴, W. JOHNSON⁶, C. NIELSEN⁷

¹V.A. & U. of Texas Health Science Center, San Antonio, TX; ²Wilford Hall Medical Ctr, San Antonio, TX; ³Fast Track Drugs & Biologics, Potomac, MD;

⁴Coley Pharmaceutical Group, Wellesley, MA; ⁵Emergent BioSolutions, Gaithersburg, MD;

⁶USAMRIID, Ft. Detrick, MD; ⁷Consultant to DARPA, Arlington, VA


ROLE OF PA IN ANTHRAX PATHOGENESIS




CPG 7909 (VAXIMMUNE)



5' – TCGTCGTTTTGTCTCGTTTTGTCCGT – 3'




MECHANISM OF ACTION OF CPG 7909



ANTHRAX VACCINE ADSORBED PLUS CPG 7909 CLINICAL STUDY DESIGN

Phase I/II proof-of-concept, double blind, randomized, controlled trial in healthy volunteers




- Healthy volunteers, age 18-45 years, male and female, at 2 study sites
- Subjects received study treatment on days 0, 14 and 28
- Subjects assessed for anti-PA and anti-TNA antibody response on days 7, 10, 14, 16, 21, 24, 28, 30, 35, 42, 49, 56.
- Safety assessed before immunization and at least weekly in clinic through day 56 and at month 6, reviewed subject-maintained diaries

Demographic Characteristics of All Randomized Subjects

Characteristic	AVA alone	CPG 7909 alone	AVA + CPG 7909	All 3 Groups
Gender – n	22	23	24	69
Male, n (%)	11 (50.0)	12 (52.2)	12 (50.0)	34 (49.3)
Age (at consent) – n	22	23	24	69
Mean (SD)	25.8 (5.8)	27.5 (4.7)	29.0 (6.7)	27.5 (5.9)
Median	24.0	26.0	27.5	26.0
Race – n (%)	22	23	24	69
White	18 (81.8)	18 (78.3)	18 (75.0)	54 (78.3)
Black/African	0 (0)	0 (0)	1 (4.2)	1 (1.5)
Amer.	1 (4.6)	1 (4.4)	4 (16.7)	6 (8.7)
Asian	0 (0)	1 (4.4)	0 (0)	1 (1.5)
Not specified	3 (13.6)	3 (13.0)	1 (4.1)	7 (10.1)
Others ^a				
Ethnicity – n (%)	22	23	24	69
Non-Hispanic	17 (77.3)	15 (65.2)	21 (87.5)	53 (76.8)
Hispanic or Latino	5 (22.7)	8 (34.8)	3 (34.8)	16 (23.2)

^aOthers include those who reported a race that was not listed on the demographics case report form or who reported multiple races. Races not represented were excluded from the table listing.

DISPOSITION OF TRIAL SUBJECTS



SUMMARY OF EFFICACY



Mean peak antibody concentration was 6.3-fold (anti-PA) and 8.8-fold (TNA) greater in the AVA plus CPG 7909 group, both $p < 0.001$

The maximum anti-PA concentration (220 μ g/mL) achieved in the AVA group (median time 42.5 days) was achieved 21 days earlier in the AVA plus CPG 7909 group ($p < 0.001$)


The maximum TNA concentration (159 μ g/mL) achieved in the AVA group (median time 46 days) was achieved 24 days earlier in the AVA plus CPG 7909 group ($p < 0.001$)

12 of 22 (55%) of subjects in the AVA plus CPG 7909 group were seropositive for anti-PA after a single immunization (day 14) compared to 2 of 18 (11%) of subjects in the AVA alone group


Significant difference in the geometric mean anti-PA antibody concentrations between the AVA and AVA plus CPG 7909 arms first detected at Day 14 ($p < 0.001$)

At peak response, 22/22 (100%) subjects in the AVA plus CPG 7909 group achieve an anti-PA concentration of $\geq 220\mu$ g/mL compared to only 11/18 (61%) of subjects in the AVA alone group

RESPONSE TO ANTRAX PA (ELISA)




TNA RESPONSE



RATE OF SEROCONVERSION




Proportion of Subjects with Anti-PA IgG Response



SAFETY



Local and Systemic Adverse Events*



* Toxicity Grading Scale for Healthy Adult and Adolescent Volunteers
Enrolled in Preventive Vaccine Clinical Trials – FDA, April 2005

LABORATORY FINDINGS

- No clinically significant changes in laboratory analyte levels were observed.
- Grade one leukopenia was seen in all study arms.
- Hypokalemia was reported in 50%, 43.5%, and 62.5 % of participants in the AVA, CPG 7909, and AVA + CPG 7909 groups, respectively, suggesting that potassium levels may have been affected by the combination treatment.

CONCLUSIONS



Safety

AVA plus CPG 7909 was reasonably well tolerated

The local injection site reactions and systemic symptoms were expected and were the most common adverse events

There was a trend to greater frequency and severity of adverse events in the AVA plus CPG 7909 group compared to the AVA alone and CPG 7909 alone groups but this was not statistically significant

Immunogenicity

AVA plus CPG 7909 elicited a heightened (6 to 8-fold increase) and accelerated (21 to 24 days) antibody response ($p < 0.001$) compared to AVA alone

AVA plus CPG 7909 elicited a positive anti-PA antibody response in >50% of subjects within 14 days of a single immunization